

SCIENTIFIC REPORTS



OPEN

Root-associated fungi of *Vaccinium carlesii* in subtropical forests of China: intra- and inter-annual variability and impacts of human disturbances

Received: 17 September 2015

Accepted: 15 February 2016

Published: 01 March 2016

Yanhua Zhang¹, Jian Ni¹, Fangping Tang¹, Kequan Pei², Yiqi Luo³, Lifen Jiang³, Lifu Sun^{1,3} & Yu Liang²

Ericoid mycorrhiza (ERM) are expected to facilitate establishment of ericaceous plants in harsh habitats. However, diversity and driving factors of the root-associated fungi of ericaceous plants are poorly understood. In this study, hair-root samples of *Vaccinium carlesii* were taken from four forest types: old growth forests (OGF), secondary forests with once or twice cutting (SEC I and SEC II), and *Cunninghamia lanceolata* plantation (PLF). Fungal communities were determined using high-throughput sequencing, and impacts of human disturbances and the intra- and inter-annual variability of root-associated fungal community were evaluated. Diverse fungal taxa were observed and our results showed that (1) Intra- and inter-annual changes in root-associated fungal community were found, and the Basidiomycota to Ascomycota ratio was related to mean temperature of the sampling month; (2) Human disturbances significantly affected structure of root-associated fungal community of *V. carlesii*, and two secondary forest types were similar in root-associated fungal community and were closer to that of the old growth forest; (3) Plant community composition, edaphic parameters, and geographic factors significantly affected root-associated fungal communities of *V. carlesii*. These results may be helpful in better understanding the maintenance mechanisms of fungal diversity associated with hair roots of ERM plants under human disturbances.

Plants of Ericaceae distribute all over the world, and are common species especially in habitats of heathlands, tundra, and forests¹. Plant species from Ericaceae could form ericoid mycorrhiza (ERM), which are expected to help them to establish successfully in habitats with low nutrients, acid soils, cold or drought stresses, and heavy metal pollutions^{2–6}. However, mycorrhizal and other fungi associated with their hosts were varied along the environment and temporal heterogeneity^{7–12}.

Significant seasonal dynamics have been observed for soil fungi^{13–15}, ectomycorrhizal (ECM) fungi^{16,17}, and arbuscular mycorrhizal (AM) fungi^{18,19} in different ecosystems. Seasonal changes of ERM fungi were also found in typical Mediterranean climate regions of Australia^{20–22}. Superiority of hair roots with ERM and their endophytes were observed in colder and more humid months²¹, and fungal richness and phylogenetic diversity was also found having the same trends under similar conditions²³. Lentendu *et al.*²⁴ suggested that soil moisture may be an important factor in determining the seasonal patterns of ERM fungi. Seasonal changes in environmental factors such as temperature, precipitation, and nutrient availability may be related to seasonal variations of fungal communities²⁴.

Fungal community changes have been observed not only in seasonality but also in inter-annual variations²⁵. These inter-annual variations may be related to both succession of plant communities and inter-annual climatic factors^{7,10,11,15,26–28}. de Román and de Miguel¹⁶ detected that percentage of ECM colonization significantly

¹College of Life Sciences, Shaoxing University, Shaoxing, China. ²State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, China. ³Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK 73019, USA. Correspondence and requests for materials should be addressed to L.S. (email: sunlifu@usx.edu.cn) or Y.Liang (email: coolrain@ibcas.ac.cn)




Figure 1. Rarefaction curves of fungal OTUs (by Chao1 estimates) in roots of *V. carlesii* in forests with different human disturbances, i.e. old growth forest (OGF), secondary forest I (SECI), secondary forest II (SECII), and plantation (PLF).

increased over 3-years period post fire, and long-term succession in microbial communities were also found in some chronosequence studies along successional gradients^{7,10,11,27,28}. Inter-annual temporal dynamics were also observed by Cotton *et al.*²⁶ in a 5-years study on AM fungal communities, in which 42% variations in AM fungal communities could be explained by the sampling year. However, the fluctuations of ERM fungal communities were less known.

Natural succession, land use changes and land management may change microclimate, edaphic and biotic factors, which would further influence diversity and composition of root-associated fungal communities^{7–12,29}. For example, with the increase of stand ages, not only fungal communities but mycorrhizal types were distinct during succession^{27,30}. Different mycorrhizal and soil fungi, such as AMF in a tropical dry ecosystem (primary forests, secondary forests and pastures) in Mexico³¹, ERM fungi in peatland sites (bog, rough grazing and forest plantation) in Ireland¹², and soil fungi in southeast Asian tropical forests (original forests, secondary forests and oil palm agriculture)³², were investigated and the results suggested that human-activity-induced changes in plant community and abiotic environments were very important in shaping fungal community compositions at the landscape scale.

To evaluate the impacts of human disturbances on root-associated fungal communities of ericaceous plants, four forest types with different human disturbances were selected: old growth forests (OGF), secondary forests with once (SEC I) and twice (SEC II) disturbances, and *Cunninghamia lanceolata* plantations (PLF). Hair roots of *V. carlesii*, a common understory ericaceous plant species in subtropical forests of China, were collected in three years (2012–2014) and root-associated fungal communities were determined using high-throughput sequencing. Fungal diversity and community composition were compared between seasons, years and different human disturbances. Three hypothesis were proposed: (1) fungal community composition has intra- and inter-annual variations, and different fungal taxa may have different temporal patterns; (2) fungal composition and community dynamic associated with hair roots of *V. carlesii* would be influenced by different human disturbances, the stronger disturbance, the larger effects on fungi; (3) plant community composition, edaphic parameters, and geographic factors may affect the composition of fungal community in roots of *V. carlesii*.

Results

Changes of *Vaccinium carlesii* population and environmental factors along with disturbance gradient.

As a common understory species in subtropical forests, *V. carlesii* was found in all four forest types along the disturbance gradient, e.g., old growth forests (OGF, without human disturbance at least 100 years), secondary forests with once clear-cut (SECI, about 50 years ago) and twice cut (SECII, clear-cut about 50 years ago, selected cut about 20 years ago), and *Cunninghamia lanceolata* plantation (PLF, clear-cut and planted *Cunninghamia lanceolata* about 20 years ago) (Fig. S1). Diameter at breast height (DBH) was much higher in OGF and individual density of *V. carlesii* was significantly higher in two secondary forests than those of OGF and PLF (Fig. S1). Soil properties in all forests were shown in Table S1. Soil pH was not significantly different in all forests, ranging from 4.73 to 4.77, a typical acid soil. Soil nutrient analysis showed that soil organic carbon (SOC), soil total nitrogen (STN), NO_3^- -N and available phosphorus (AP) were, but soil total phosphorus (STP) and NH_4^+ -N were not significantly different along the disturbance gradient, indicating that more available nutrients appeared in OGF, and no different nutrient supply between disturbed forest types.

Fungal diversity in hair roots of *V. carlesii*.

Rarefaction curves of fungal OTUs in roots of *V. carlesii* were shown in Fig. 1. While the mean number of OTUs in SECI was slightly lower, no significant differences were found between four forest types. There were 5595 OTUs in hair roots of *V. carlesii* in all forest types, including Ascomycota, Basidiomycota, Zygomycota, Chytridiomycota, and Glomeromycota (Fig. 2). Ascomycota and Basidiomycota were two dominant phyla in all forest types, and proportions of Zygomycota, Chytridiomycota, and Glomeromycota were only 1.14%, 0.08%, and 0.03%, respectively. Common classes included Leotiomycetes, Eurotiomycetes, Dothideomycetes, Sordariomycetes from Ascomycota and Agaricomycetes from Basidiomycota. Dominant fungal orders (>5%) included typical ERM orders such as Helotiales (21.9%), Sebaciniales (8.9%), and Chaetothyriales (6.1%), as well as typical ECM orders such as Thelephorales (9.0%) and Russulales (8.9%).




Figure 2. Proportions of fungal phyla and common fungal classes (>1%) within Ascomycota and Basidiomycota in different forest types, seasons, and years (only samples taken in spring were compared).

Venn diagrams showed the number of specific and shared OTUs of *V. carlesii* by seasons (Fig. 3B), years (Fig. 3C) and different human disturbances (Fig. 3A). Figure 3(B) showed that 494 OTUs was shared among four seasons in 2012, which was 10.36% of total 4768 OTUs. The seasonal specific OTUs in spring, summer, autumn and winter of 2012 were 988, 703, 612 and 656, accounting for 20.72%, 14.74%, 12.84% and 13.76% of total OTUs, respectively. Both specific and total OTU number in spring was much higher than those in the other three seasons.

In Fig. 3(C), of the total 3726 OTUs that were observed in spring of three years, only 526 OTUs were found in all three year, accounting for 87.1% of total reads in each sample. The specific OTUs occurred only in 2012 were much more than those in 2013 and 2014.

Total 2518, 2170, 2191, and 2435 OTUs were found in OGF, SECI, SECII, and PLF (Fig. 3A), of which 962, 653, 814, and 1035 OTUs were specific to these four forest types, respectively. 502 OTUs were shared by all forest types, accounting for 8.9% of the total OTU number. When excluding 502 OTUs shared by all four forest types, only 73 OTUs were found in both PLF and two secondary forests, which was much less than the shared OTUs between OGF and the secondary forests (347 OTUs).

Fungal community composition varied with season and year and responses to human disturbance. Principal Component Analysis (PCA) results on community structure of root-associated fungi of *V. carlesii* in different forests sampled in different seasons of 2012 and spring of 2013 and 2014 were shown in Figs 4 and S2. Four seasons were not well separated in 2012 and root-associated fungal communities in spring of 2013 and 2014 were much distinct to those of 2012. The results of dominant phyla and classes showed that cold (spring and winter) and warm (summer and autumn) seasons had different compositions at high taxa level (Fig. 2B). The ratio of Basidiomycota to Ascomycota in root-associated fungal communities was significantly higher in summer and autumn and significantly correlated with mean air temperature of the sampling month ($r = 0.698$, $P < 0.001$, Fig. 5).

The results of PCA showed that fungal community structures of two secondary forests were more similar to old growth forests than to plantations for all six sampling dates (Figs 4 and S3).

Indicator fungal species for seasons, years and forest types. There were 27 OTUs that showed significant preference to a specific season (Table 1), including 15, 2, 8, and 2 indicator species for spring, summer, autumn and winter, respectively. Among the indicator species for seasons, Helotiales 3, Helotiales 4, Helotiales 5, Dermateaceae 1, Herpotrichiellaceae 4 and Herpotrichiellaceae 5 are putative ERM fungi^{1,22,33–35}; and Thelephoraceae (Thelephoraceae 1 and Thelephoraceae 2) are usually considered as members of ECM fungal family³⁶.




Figure 3. Venn diagram showing specific and shared OTUs of different forest types (A), seasons (B), and years (C) (only samples taken in spring were compared).




Figure 4. Principal Component Analysis (PCA) of root-associated fungi of *V. carlesii*. Forest types are shown as different shapes (OGF circle, SEC I square, SEC II triangle, PLF diamond), and seasons are shown as different colors (Spring green, Summer red, Autumn brown, Winter white).

There were 3, 6, and 14 indicator fungal OTUs for three sampling years (Table 2). In March of 2013, all the indicator species are putative ERM fungi, such as *Cryptosporiopsis ericae*^{34,37}, *Oidiendron maius*^{22,37,38}, Myxotrichaceae 2 and three species of Dermateaceae (Helotiales). Indicators for March of 2014 were diverse, including putative ERM fungi (Helotiales 6, Helotiales 7, Helotiales 8, Hypocreales 1^{35,39,40}, and Sebaciales 1), putative ECM fungi (*Tomentella* sp1 and *Tomentellopsis* sp1), as well as saprotrophic fungi (Malasseziales 1, *Penicillium herquei*, *Penicillium* sp3, *Penicillium* sp6 and *Penicillium* sp7).

There were 22 OTUs that exhibited significant preference to human disturbances (Table 3). The numbers of indicator species for OGF, SECI, SECII, and PLF were 7, 0, 2, and 13, respectively. The low number of indicator species in two secondary forests may be related to their similar abiotic and biotic factors to the old growth forest, and much higher number of indicator species in PLF may be due to the distinct abiotic and biotic factors in plantations. Four indicative fungal species in OGF belong to putative ERM fungi, i.e. Myxotrichaceae 1 and three OTUs of Herpotrichiellaceae (Herpotrichiellaceae 1, Herpotrichiellaceae 2 and *Cladophialophora chaetospora*).




Figure 5. Relationship between mean air temperature of the sampling month and ratio of Basidiomycota to Ascomycota in root-associated fungal communities. Sampling season □ winter, ■ spring, ■ summer, ■ autumn.

Index	Identified name	Group	P
1	Dermateaceae 1	Spring	0.019
2	Helotiales 2	Spring	0.025
3	Helotiales 3	Spring	0.016
4	Helotiales 4	Spring	0.003
5	Helotiales 5	Spring	0.022
6	Herpotrichiellaceae 4	Spring	0.039
7	Leotiomycetes 1	Spring	0.011
8	<i>Mortierella</i> sp1	Spring	0.008
9	<i>Parmelia</i> sp1	Spring	0.021
10	<i>Penicillium</i> sp3	Spring	0.043
11	<i>Penicillium spinulosum</i>	Spring	0.048
12	<i>Phialocephala fortinii</i>	Spring	0.044
13	<i>Pochonia bulbilosa</i>	Spring	0.013
14	Thelephoraceae 1	Spring	0.011
15	unidentified fungus 2	Spring	0.011
16	Eurotiomycetes 1	Summer	0.021
17	<i>Trichoderma</i> sp1	Summer	0.02
18	Ascomycota 4	Autumn	0.043
19	Ascomycota 5	Autumn	0.002
20	Agaricomycetes 1	Autumn	0.021
21	Basidiomycota 1	Autumn	0.031
22	Herpotrichiellaceae 5	Autumn	0.001
23	<i>Paecilomyces</i> sp1	Autumn	0.018
24	<i>Penicillium</i> sp4	Autumn	0.005
25	<i>Penicillium</i> sp5	Autumn	0.005
26	Thelephoraceae 2	Winter	0.049
27	Trechisporales 1	Winter	0.049

Table 1. Indicator fungal species associated with hair roots of *V. carlesii* in four sampling seasons.

Of the indicator species for PLF, *Sebacina* sp1, *Sebacina* sp2, Herpotrichiellaceae 3 and Chaetothyriales 1 are from common ERM fungal orders, and *Diaporthe* sp1 and *Pestalotiopsis* sp1 are considered as plant pathogens.

Factors affect fungal community associated with *V. carlesii* hair roots. Our results also showed that plant community, soil parameters, and geographic factors had significant effects on root-associated fungal community of *V. carlesii* (Table 4). For plant community, PC2, PC3 and species richness showed significant effects. The plant species that had significant contribution on PC2 and PC3 included the commercial species in PLF, i.e., *Cunninghamia lanceolata*, some dominant ECM plant species in subtropical forest, i.e., *Castanopsis eyrei* and *Lithocarpus glaber*, as well as some common ERM plant species, i.e., *R. ovatum*, *R. latoucheae* and *V. carlesii*.

Index	Identified name	Group	P
1	Eurotiomycetes 2	2012	0.0016
2	unidentified fungus 3	2012	0.0006
3	unidentified fungus 4	2012	0.0074
4	<i>Cryptosporiopsis ericae</i>	2013	0.0052
5	Dermateaceae 2	2013	0.0004
6	Dermateaceae 3	2013	0.0037
7	Dermateaceae 4	2013	0.0047
8	Myxotrichaceae 2	2013	0.0005
9	Oidiodendron maius	2013	<0.0001
11	Debaryomyces sp1	2014	0.0007
10	Helotiales 6	2014	0.0065
12	Helotiales 7	2014	0.0083
13	Helotiales 8	2014	0.0022
14	Hypocreales 1	2014	0.0077
15	Malasseziales 1	2014	0.0001
16	<i>Penicillium herquei</i>	2014	0.0068
17	<i>Penicillium</i> sp3	2014	0.0023
18	<i>Penicillium</i> sp6	2014	0.0001
19	<i>Penicillium</i> sp7	2014	0.0018
20	Sebacinales 1	2014	0.0059
21	<i>Tomentella</i> sp1	2014	0.0044
22	<i>Tomentellopsis</i> sp1	2014	0.0015
23	unidentified fungus 5	2014	0.0096

Table 2. Indicator fungal species associated with hair roots of *Vaccinium carlesii* for spring of different sampling years.

Index	Identified name	Group	P
1	Ascomycota 1	OGF	0.0002
2	Myxotrichaceae 1	OGF	0.0001
3	Herpotrichiellaceae 1	OGF	0.0001
4	Herpotrichiellaceae 2	OGF	0.0018
5	<i>Cladophialophora chaetospira</i>	OGF	0.0002
6	<i>Penicillium</i> sp1	OGF	0.0020
7	unidentified fungus 1	OGF	0.0025
8	Sebacinaeae 1	SECII	0.0009
9	Ascomycota 2	SECII	0.0017
10	Helotiales 1	PLF	0.0008
11	<i>Penicillium</i> sp2	PLF	0.0015
12	<i>Diaporthe</i> sp1	PLF	0.0048
13	<i>Trichoderma</i> sp1	PLF	0.0028
14	<i>Neonectria</i> sp1	PLF	0.0055
15	<i>Pestalotiopsis</i> sp1	PLF	0.0099
16	Ascomycota 3	PLF	0.0014
17	<i>Clitopilus prunulus</i>	PLF	0.0036
18	Sebacina sp1	PLF	<0.0001
19	Sebacina sp2	PLF	<0.0001
20	Herpotrichiellaceae 3	PLF	<0.0001
21	Sordariomycetes 1	PLF	0.0024
22	Chaetothyriales 1	PLF	0.0044

Table 3. Indicator fungal species associated with hair roots of *V. carlesii* in forests with different human disturbances.

These results indicated that both dominant trees and common ERM neighbors may affect fungal community of *V. carlesii*. For soil parameters, soil organic carbon (SOC), soil total nitrogen (STN), soil total phosphorus (STP), and ammonium nitrogen ($\text{NH}_4^+\text{-N}$) had significant effects on fungal community of *V. carlesii*. Longitude and

	RDA1	RDA2	r ²	P
Plant community				
PC1	-0.774	0.633	0.005	0.845
PC2	0.980	0.201	0.188	<0.001
PC3	-0.443	0.896	0.186	0.003
Species richness	0.830	-0.558	0.082	0.050
TABH	-0.957	0.292	0.065	0.094
Edaphic parameters				
SOC	0.829	-0.560	0.129	0.012
STN	0.839	-0.544	0.114	0.019
STP	0.303	-0.953	0.100	0.025
NH ₄ ⁺ -N	0.779	-0.627	0.142	0.006
NO ₃ ⁻ -N	0.729	0.684	0.060	0.118
AP	0.865	-0.501	0.029	0.362
pH	-0.589	-0.808	0.014	0.628
Geographic factors				
Longitude	-0.667	-0.745	0.154	0.003
Latitude	0.507	0.862	0.215	<0.001
Altitude	-0.392	0.920	0.060	0.116

Table 4. Correlations of microbial community composition with plant community, edaphic and geographic factors. Abbr.s: TABH, total area at breast height; PC1, PC2 and PC3, the first three principal components of plant community; SOC: soil organic carbon, STN: soil total nitrogen content, STP: soil total phosphorus content, NH₄⁺-N: ammonium nitrogen, NO₃⁻-N: nitrate nitrogen, AP: Available phosphorus.

	df	SS	MS	F	P
Forest type	3	2.650	0.883	3.291	0.001
Year	2	2.769	1.385	5.159	0.001
Season	3	0.922	0.307	1.145	0.204
Longitude	1	0.441	0.441	1.645	0.046
Latitude	1	0.560	0.560	2.086	0.007
Altitude	1	0.369	0.369	1.376	0.096
Residuals	60	16.103	0.268	0.676	

Table 5. Adonis results showing effects of different factors on root-associated fungal community of *Vaccinium carlesii*.

latitude are significant geographic factors, showing that geographic distribution of plots may have significant effects on fungal community of *V. carlesii*.

The Adonis results showed that forest type, year, longitude, and latitude had significant effects on fungal community of *V. carlesii* (Table 5). The effects of season and altitude were not significant.

Discussion

Diversity of total fungal OTUs associated with *V. carlesii* hair roots. Diverse fungal species were observed in hair roots of *V. carlesii* in the present study. These fungal species include typical ERM fungal orders such as Helotiales and Sebaciales, which have been observed in roots of many ericaceous plant species^{1,33,34,37,38,41,42}. Some fungal families, such as Myxotrichaceae^{37,39}, Herpotrichiellaceae^{41,43}, Dermateaceae^{1,22,34,35}, and Sebacinaceae^{6,43,44} considered as ERM fungi, were frequently found in our results. Some putative ERM fungal taxa were detected in our study, such as *Lachnum*¹, *Scytalidium*⁴⁰, *Meliniomyces*^{1,29}, *Gliocladium*³⁹, *Cryptosporiopsis ericea*¹, *Oridiodendron maius*^{2,22,37,39,45} and *Rhizoscyphus ericae*^{22,45,46}. The latter two fungal species which were frequently cultured from ericaceous roots as typical ERM fungi, was less in our results, and coincidence with the results from Allen *et al.*⁴³ and Bougoure and Cairney⁴⁷ research, suggesting that these fungi were not dominants of ERM fungal communities in open ecosystems.

Some typical ECM fungal genera were also found in roots of *V. carlesii*, for example *Russula*, *Tomentella*, *Rhizopogon*, *Thelephora*, *Cenococcum*. Those fungal genera were also observed in roots of some other ericaceous plants^{36,42,43}, indicating that they may form symbiotic structures with both ERM and ECM host plants. In subtropical forests of China, the dominant tree species of canopy are usually ECMF hosts (e.g. Fagaceae) and Ericaceae plants are dominant in the shrub layer. Common mycorrhizal networks formed between ECM and ERM hosts may be essential for species coexistence and ecosystem functioning in subtropical forests.

Besides of mycorrhizal fungi, DSE (e.g., *Phialocephala fortinii*^{1,41,43,47,48}), saprobes^{34,35,42}, pathogens^{1,42} and unidentified root-associated fungi were also present in hair roots of *V. carlesii*. DSE was found frequently co-exist

in ericaceous roots^{30,32,49} in heathlands, forests and alpine ecosystems⁵⁰ and showed stronger resistant to adverse conditions such as drought, repeated freezing and thawing⁵¹. Ecological functions of saprobes (such as *Penicillium* spp.) observed in our investigation are still unknown, while it has been documented that some typical saprophytic fungal genera were also observed in roots of ERM plants in previous studies, e.g., *Acremonium*⁴⁰, *Capronia*^{42,43}, *Myrothecium*⁴³. One possible reason for the occurrences of saprobes is that these fungi may act as endophytes at some stages of their lifecycle and saprobes in soil at other stages. It has been found that putative ERM fungi (e.g. *Oidiodendron maius*^{52,53} and *R. ericae*^{53,54}) could live as saprotrophs for a long time when their hosts were absent⁵⁵.

Compared with traditional culture-based, isolated and morphological identified approaches, application of high-throughput sequencing makes a big progress in the studies of mycorrhizal and soil fungal diversity and their communities^{23,24,56–60}, because it can generate unprecedented numbers of sequences, and even very rare and low-abundance organisms^{61–63} can be detected. We observed 5595 fungal OTUs in our study sites, while in researches using on isolation and culture or local database based ITS-RFLP (Internal Transcribed Spacer-Restriction Fragment Length Polymorphism), Zhang *et al.*³⁷, Tian *et al.*³⁹, Sun *et al.*⁴⁴ found only 17, 12 and 35 fungal taxa, respectively. Helotiales and Sebaciniales were usually found in the previous studies in ericaceous plants⁴⁴. Our results using high-throughput sequencing not only found these fungal taxa, but also diverse fungal taxa in roots of *V. carlesii*. Diverse fungal taxa were also observed when studying other mycorrhizal fungal community using high-throughput sequencing. Buscardo *et al.*⁶⁰ supposed that the number of ECMF taxa was eight-fold higher by using 454 pyrosequencing than by DGGE (Denaturing Gradient Gel Electrophoresis), and Oja Jane *et al.*⁵⁹ obtained 5805 OTUs in orchid mycorrhizal symbionts by 454 pyrosequencing.

Intra- and inter- annual dynamics of root-associated fungal communities. Seasonal changes of dominant fungal taxa in roots of *V. carlesii* have been found in the present study (Fig. 2B) and the ratio of Basidiomycota to Ascomycota is significantly correlated with monthly mean temperature (Fig. 5). In another study conducted in the typical Mediterranean climate regions of Australia, researchers also found significant seasonal variations of ERM fungi, in which ERM colonization and diversity were higher in colder and more humid winter^{20,21,23}. Soil moisture has been suggested as a key factor in determining seasonal changes of ERM communities²⁴. However, in the subtropical region of our study, the highest precipitation usually occurs from June to August and water might not always be a limiting factor in determining seasonal patterns of fungal community in roots of *V. carlesii*. The total and specific number of OTUs occurred in spring were more than in three other seasons, implying that the colder and more humid climate in our investigation region might be benefit to most fungi development.

The seasonal changes of common fungal taxa and the ratio of Basidiomycota to Ascomycota also imply that different fungal taxa may respond differently to seasonal environmental changes and interactions between fungal taxa may also change within a year. Seasonality has been reported in AM and ECM fungal communities^{16,17,64,65} and soil fungi⁶⁶, and seasonal changes may be initialized from ecological factors such as temperature, moisture, vegetation-soil interactions and substrate availability changes.

OTU numbers in spring of 2012 were extremely higher than that in 2013 and 2014 (Fig. 3C) and the results of PCA showed significant difference in root associated fungal community between years (Fig. 4). When comparing the samples collected in spring from 2012 to 2014, obvious fluctuations of fungal communities were observed (in Figs 4 and S2). Most of Zhejiang province including our study site experienced an extremely drought during the summer of 2013, which could affect fungal of 2014 spring. The distinct fungal community in spring of three years and more indicator species for the year 2014 could be partially due to the inter-annual shifts of climatic factors. The mechanisms underlying inter-annual shifts of root-associated fungal community may be quite complex, and changes in plant community, edaphic factors, and climatic parameters may be involved. For example, roots development and mycorrhizal infection may be affected by moisture regimes, which depend on the rainfall variations related with the El Niño or La Nina years in Australia^{20,43}. Cairney & Ashford²⁰ found that in El Niño year with summer drought, ERM colonization and amount of hair roots are positively correlated with soil moisture, while in La Nina year without summer drought, temperature was the determinant factor for ERM structures²⁰. Therefore, limited factors on hair roots and ERM fungi development were different by seasons and years. Sometimes sampling time can also strongly affect the microbial community compositions⁶⁷.

Root associated fungal community along human disturbance gradients. All the forest types once covered by the same vegetation of subtropical forests, with dominant plant families of Fagaceae, Theaceae, Lauraceae, Pinaceae, Taxodiaceae, etc. Human disturbance extremely altered dominant species composition of plants, especially in *Cunninghamia lanceolata* plantation (Table S2).

Our results showed that human disturbances, especially the *C. lanceolata* plantation significantly changed community structure of root associated fungi of *V. carlesii*. The number of total and specific OTUs in PLF was much higher than those in two secondary forests with more DSE and pathogen species, and both of these fungal types were found often present in earlier stages of succession^{27,68,69} with less mycorrhizal fungi³⁰. More proportion of ERM fungi co-existed in the fungal communities in OGF probably due to the colonization rate of mycorrhizal fungi would increase along with the soil development years^{21,70}. Compared with plantations, secondary forests have more similar root-associated fungal communities with each other and closer to the old growth forest, which may be attributed to the relatively similar accompanying plant compositions and similar abiotic environmental parameters between old growth forests and secondary forests. The distinct root-associated fungal community of *V. carlesii* in PLF might be due to (1) the different mycorrhizal status of dominant plants between PLF and other forest types, since *C. lanceolata* is AM plants⁷¹ while the dominant plant family (Fagaceae) is ectomycorrhizal; (2) the differences in soil parameters between PLF and other three forest types; and (3) the spatial aggregation and isolation. Our results also showed that plant community, soil parameters, and geographic factors had significant effects on root-associated fungal community of *V. carlesii* (Table 4). Results from some previous studies also

suggested that old forests and secondary forests were more similar in fungal communities though the disturbance patterns were different^{24,32,72}, and they would be more and more similar in soils over time^{7,10,11}.

Fungal diversity and its dynamic changes have been investigated after plant composition and abiotic factors varied^{12,13,70}. Environmental changes make the same hosts have different fungal communities, for example, Bougoure *et al.*⁷³ examined root-associated fungal communities of *Calluna vulgaris* along a heath to forest gradient in Scotland and found significant differences; and Hazard *et al.*¹² also found that diversity of fungi associated with hair roots of *Vaccinium macrocarpon* in peatland of Ireland was affected by three different land use (bog, rough grazing and forest plantation). Human disturbance happened frequently in forest ecosystems, however, in some cases, even after over fifty years of conversion from land use^{8,9}, or more than one century succession⁷⁰, significant difference between older and earlier forest ecosystems would still be observed in microbial community structure. The effects of disturbance on the species and richness of soil and mycorrhizal fungi were also found in other studies on ECM, AM and soil fungal communities^{29,70,74}. The responses of root-associated fungi to human disturbance gradients may depend much on the host plants, the ecosystems selected and the extent of disturbance gradients.

Conclusions

Diverse fungal OTUs have been observed in hair-roots of *Vaccinium carlesii* using high-throughput sequencing, including putative ERM fungi, ECM fungi, common DSE, saprobes and pathogens. Dominant phyla are Ascomycota and Basidiomycota, and common classes are Leotiomycetes, Eurotiomycetes, Dothideomycetes, Sordariomycetes and Agaricomycetes. Intra- and inter- annual variability in root-associated fungal community of *V. carlesii* have been observed in the present study, and the ratio of Basidiomycota to Ascomycota is related to monthly mean temperature of the sampling month. Significant differences were found between different forest types along the disturbance gradient, and root-associated fungal communities of *V. carlesii* in two secondary forest types are similar with each other and are closer to that in old growth forests. Factors affecting root-associated fungal communities of *V. carlesii* include plant community composition, edaphic parameters, and geographic factors.

Materials and Methods

Study sites. The study site is located at Gutianshan National Nature Reserve (GNNR), Zhejiang Province in East China (29°10′19.4″N–29°17′41.4″N, 118°03′49.7″E–118°11′12.2″E). Annual mean temperature is 15.3 °C and annual precipitation ranges from 1793 to 1960 mm. Subtropical red soil with granite or deeply weathered granite as parent rock is the dominant soil type⁷⁵. The typical vegetation in this region is subtropical evergreen broad-leaved forest⁷⁶, with *Castanopsis eyrei* and *Schima superba* being the dominant canopy species, and *Vaccinium carlesii* is one of the common understory shrubs with abundant individuals in this area (Table S2).

Four types of forests with different disturbance history in the GNNR were studied in the present study: old growth forests (OGF), secondary forests with once cutting (SEC I), secondary forests with twice cutting (SEC II), and *Cunninghamia lanceolata* plantation (PLF). Within each type of forests, three 1-ha (100 m × 100 m) plots were randomly selected. SEC I was clearly cut about 50 years ago, while SEC II was clearly cut about 50 years ago and then selectively cut about 20 years ago. PLF was planted about 20 year ago after clear cutting of secondary forests. Stands in both types of secondary forests and plantations have been undergoing natural recovery since last anthropogenic disturbances. OGF is undisturbed forests that did not experience tree-felling during the last 100 years and is generally located at the core zone of GNNR⁷⁷.

Sampling procedure. Hair roots of four individuals of *V. carlesii* were sampled in March, June, September, December of 2012 and March of 2013 and 2014 from each plot. In total, root samples from 288 individuals were collected (4 samples × 12 plots × 6 sampling times). In detail, the terminal portion of the finer roots (typical ericaceous “hair roots”) was retrieved from soils at four directions around the trunk of each *V. carlesii* individual. The roots were washed carefully after 1 h soaking in sterile water. Hair roots were then cut into 1 cm segments and 20 hair root segments were selected randomly from each *V. carlesii* individual sample. Each root segment was put into a centrifuge tube and preserved in 70% alcohol at –70 °C before DNA extraction. A 200 g soil sample was taken from the top 10 cm of soil adjacent to each plant sampled for elemental analyses.

DNA extraction. DNA was extracted from hair roots of *V. carlesii* following the protocol of DNA secure Plant Kit (TIANGEN Biotech Co. Ltd), with slight modifications. Hair root segments were put into a sterile centrifuge tube containing 20 µl 2× CTAB extraction buffer solution and ground with a plastic pestle on ice. Samples were warmed at 65 °C for 1 h in 630 µl aliquot of 2× CTAB extraction buffer solution, and then shaken for 10 min. Aliquot of chloroform/isoamyl alcohol (24:1) was added followed by twice centrifugation at 13,201 × g for 8 min at room temperature. The supernatant was precipitated with 100% alcohol at 4 °C for 1 h followed by centrifugation at 17,968 × g for 8 min. DNA precipitate was washed twice using 70% ethanol, dried in a vacuum desiccator, and dissolved in 30 µl sterile ddH₂O at 4 °C. DNA samples were stored at –22 °C prior to downstream analyses.

PCR amplifications. The ITS1F-ITS2 region of fungi were amplified by PCR (95 °C for 2 min, followed by 25 cycles at 95 °C for 30 s, 55 °C for 40 s, and 72 °C for 50 s and a final extension at 72 °C for 5 min) using primers ITS1F 5′-barcode-CTTGGT CATTTAGAGGAAGTAA-3′ and ITS2 5′-GCTGCGTTCTTCATCGATGC-3′, where barcode is an eight-base sequence unique to each sample. PCR reactions were performed in triplicate 20 µl mixture containing 4 µl of 5× FastPfu Buffer, 2 µl of 2.5 mM dNTPs, 0.8 µl of each primer (5 µM), 0.4 µl of FastPfu Polymerase, and 10 ng of template DNA.

Illumina MiSeq sequencing. Amplicons were extracted from 2% agarose gels and purified using the AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, Union City, CA, US) following the manufacturer's instructions and quantified using QuantiFluor™-ST (Promega, U.S.). Purified amplicons were pooled in equimolar and paired-end sequenced (2 × 250) on an Illumina MiSeq platform adopting the standard protocols.

Bioinformatic analysis. Raw fastq files were demultiplexed and quality-filtered using QIIME (ver 1.7) with the following criteria: (i) the reads were truncated at any site receiving an average quality score <20 over a 10 bp sliding window, discarding the truncated reads that were shorter than 50 bp; (ii) exact barcode matching, 2 nucleotide mismatch in primer matching, reads containing ambiguous characters were removed; (iii) only sequences that overlapped by longer than 10 bp were assembled according to their overlap sequence. Reads that could not be assembled were discarded.

Open reference OTU picking was done with `pick_open_reference_otus.py` using the default `uclust` method and ITS 12-11 dataset (97% similarity cutoff was used, alpha release, download from web site of QIIME http://qiime.org/home_static/dataFiles.html), and singletons were removed during OTU picking. The phylogenetic affiliation of each sequence was analyzed by RDP Classifier (<http://rdp.cme.msu.edu/>) against ITS-12-11 dataset using confidence threshold of 80%. `Single_rarefaction.py` in Qiime was used to generate OTU table with even reads of 10,000 in each root sample.

Statistical analysis. One-way ANOVA (with post hoc comparisons using Duncan's test) was carried out to test the difference of individual density and mean DBH per individual of *V. carlesii* between four forest types using SPSS (ver. 16.0, SPSS Inc.). Indicator species analysis was performed using `out_category_significance.py` in Qiime to determine the indicator fungal species for different forest types, seasons, and years. "Envfit" function was used to identify the main factors influencing root-associated fungal community. Principal Components Analysis (PCA) was performed using "rda" function in the R package "vegan" and the first three components (PC1, PC2, and PC3) were used as plant parameters in "envfit". "Adonis" function in the R package "vegan" was used to evaluate the impacts of forest type, year, season, and geographic factors on root-associated fungal community of *Vaccinium carlesii*.

Ethics Statement. No specific permits were required for the described field studies. The study sites are not privately-owned or protected in any way, and the field studies did not involve endangered or protected species.

References

- Walker, J. F. *et al.* Diverse Helotiales associated with the roots of three species of Arctic Ericaceae provide no evidence for host specificity. *New Phytol.* **191**, 515–527 (2011).
- Martino, E., Franco, B., Piccoli, G., Stocchi, V. & Perotto, S. Influence of zinc ions on protein secretion in a heavy metal tolerant strain of the ericoid mycorrhizal fungus *Oidiodendron maius*. *Mol. Cell Biochem.* **231**, 179–185 (2002).
- Cairney, J. W. G. & Meharg, A. A. Ericoid mycorrhiza: a partnership that exploits harsh edaphic conditions. *Eur. J. Soil Sci.* **54**, 735–740 (2003).
- Persson, J. *et al.* Nitrogen acquisition from inorganic and organic sources by boreal forest plants in the field. *Oecologia* **137**, 252–257 (2003).
- Rains, K. C. & Bledsoe, C. S. Rapid uptake of 15N-ammonium and glycine-13C, 15N by arbuscular and ericoid mycorrhizal plants native to a Northern California coastal pygmy forest. *Soil Biol. Biochem.* **39**, 1078–1086 (2007).
- Ishida, T. A. & Nordin, A. No evidence that nitrogen enrichment affect fungal communities of *Vaccinium* roots in two contrasting boreal forest types. *Soil Biol. Biochem.* **42**, 234–243 (2010).
- Buckley, D. H. & Schmidt, T. M. Diversity and dynamics of microbial communities in soils from agro-ecosystems. *Environ. Microbiol.* **5**, 441–452 (2003).
- Fraterrigo, J. M., Turner, M. G., Pearson, S. M. & Dixon, P. Effects of past land use on spatial heterogeneity of soil nutrients in southern Appalachian forests. *Ecol. Monogr.* **75**, 215–230 (2005).
- Fraterrigo, J. M., Balser, T. C. & Turner, M. G. Microbial community variation and its relationship with nitrogen mineralization in historically altered forests. *Ecology* **87**, 570–579 (2006).
- Jangid, K., Williams, M. A., Franzluebbers, A. J., Blair, J. M. & Coleman, D. C. Development of soil microbial communities during tall grass prairie restoration. *Soil Biol. Biochem.* **42**, 302–312 (2010).
- Jangid, K., Williams, M. A., Franzluebbers, A. J., Schmidt, T. M. & Coleman, D. C. Land-use history has a stronger impact on soil microbial community than aboveground vegetation and soil properties. *Soil Biol. Biochem.* **43**, 2185–2193 (2011).
- Hazard, C., Gosling, P., Mitchell, D. T., Doohan, F. M. & Bending, G. D. Diversity of fungi associated with hair roots of ericaceous plants is affected by land use. *FEMS Microbiol. Ecol.* **87**, 586–600 (2014).
- Carney, K. M. & Matson, P. A. The influence of tropical plant diversity and composition on soil microbial communities. *Microbiol. Ecol.* **52**, 226–238 (2006).
- Buckeridge, K. M., Banerjee, S., Siciliano, S. D. & Grogan, P. The seasonal pattern of soil microbial community structure in mesic low arctic tundra. *Soil Biol. Biochem.* **65**, 338–347 (2013).
- Schadt, C. W., Martin, A. P., Lipson, D. A. & Schmidt, S. K. Seasonal dynamics of previously unknown fungal lineages in tundra soils. *Science* **301**, 1359–1361 (2003).
- de Román, M. & de Miguel, A. M. Post-fire, seasonal and annual dynamics of the ectomycorrhizal community in a *Quercus ilex* L. forest over a 3-year period. *Mycorrhiza* **15**, 471–482 (2005).
- Morgado, L. N. *et al.* Summer temperature increase has distinct effects on the ectomycorrhizal fungal communities of moist tussock and dry tundra in Arctic Alaska. *Glob. Change Biol.* **21**, 959–972 (2015).
- Soteras, F., Grilli, G., Cofré, M. N., Marro, N. & Becerra, A. Arbuscular mycorrhizal fungal composition in high montane forests with different disturbance histories in central Argentina. *Appl. Soil Ecol.* **85**, 30–37 (2015).
- Mandyam, K. & Jumpponen, A. Seasonal and temporal dynamics of arbuscular mycorrhiza and dark septate endophytic fungi in a tallgrass prairie ecosystem are minimally affected by nitrogen enrichment. *Mycorrhiza* **18**, 145–155 (2008).
- Cairney, J. W. G. & Ashford, A. E. Tansley review no. 135. Biology of mycorrhizal associations of epacrids (Ericaceae). *New Phytol.* **154**, 305–326 (2002).
- Hutton, B. J., Dixon, K. W. & Sivasithampamram, K. Effect of habitat disturbance on inoculum potential of ericoid endophytes of Western Australian heaths (Epacridaceae). *New Phytol.* **135**, 739–744 (1997).

22. Usuki, F., Abe, J. P. & Kakishima, M. Diversity of ericoid mycorrhizal fungi isolated from hair roots of *Rhododendron obtusum* var. *kaempferi* in a Japanese red pine forest. *Mycoscience* **44**, 97–102 (2003).
23. Pellissier, L. *et al.* Soil fungal communities of grasslands are environmentally structured at a regional scale in the Alps. *Mol. Ecol.* **23**, 4274–4290 (2014).
24. Lentendu, G., Zinger, L., Manel, S., Coissac, E. & Choler, P. Assessment of soil fungal diversity in different alpine tundra habitats by means of pyrosequencing. *Fungal Divers.* **49**, 113–123 (2011).
25. Peršoh, D. Plant-associated fungal communities in the light of metaomics. *Fungal Divers.* **75**, 1–25 (2015).
26. Cotton, T. E., Fitter, A. H., Miller, R. M., Dumbrell, A. J. & Helgason, T. Fungi in the future: interannual variation and effects of atmospheric change on arbuscular mycorrhizal fungal communities. *New Phytol.* **205**, 1598–1607 (2015).
27. Dickie, I. A. *et al.* Mycorrhizas and mycorrhizal fungal communities throughout ecosystem development. *Plant Soil* **367**, 11–39 (2013).
28. Welc, M., Frossard, E., Egli, S., Bünemann, E. K. & Jansa, J. Rhizosphere fungal assemblages and soil enzymatic activities in a 110-years alpine chronosequence. *Soil Biol. Biochem.* **74**, 21–30 (2014).
29. Grellet, G. A., Johnson, D., Paterson, E., Münzenberger, B. & Hüttl, R. F. Reciprocal carbon and nitrogen transfer between an ericaceous dwarf shrub and fungi isolated from *Piceirhiza bicolorata* ectomycorrhizas. *New Phytol.* **182**, 359–366 (2009).
30. Cázares, E., Trappe, J. M. & Jumpponen, A. Mycorrhiza-plant colonization patterns on a subalpine glacier forefront as a model system of primary succession. *Mycorrhiza* **15**, 405–416 (2005).
31. Gavito, M. E., Pérez-Castillo, D., González-Monterrubio, C. F., Vieyra-Hernández, T. & Martínez-Trujillo, M. High compatibility between arbuscular mycorrhizal fungal communities and seedlings of different land use types in a tropical dry ecosystem. *Mycorrhiza* **19**, 47–60 (2008).
32. McGuire, K. L. *et al.* Responses of Soil Fungi to Logging and Oil Palm Agriculture in Southeast Asian Tropical Forests. *Microb. Ecol.* **8**, 1–15 (2014).
33. Gorzelak, M. A., Hambleton, S. & Massicotte, H. B. Community structure of ericoid mycorrhizas and root-associated fungi of *Vaccinium membranaceum* across an elevation gradient in the Canadian Rocky Mountains. *Fungal Ecol.* **5**, 36–45 (2012).
34. Chambers, S. M., Curlevski, N. J. A. & Cairney, J. W. G. Ericoid mycorrhizal fungi are common root inhabitants of non-Ericaceae plants in a south-eastern Australian sclerophyll forest. *FEMS Microbiol. Ecol.* **65**, 263–270 (2008).
35. Bergero, R., Girlanda, M., Bello, F., Luppi, A. & Perotto, S. Soil persistence and biodiversity of ericoid mycorrhizal fungi in the absence of the host plant in a Mediterranean ecosystem. *Mycorrhiza* **13**, 69–75 (2003).
36. Smith, J. E., Molina, R. & Perry, D. A. Occurrence of Ectomycorrhizas on ericaceous and coniferous seedlings grown in soils from the Oregon coast range. *New Phytol.* **129**, 73–81 (1995).
37. Zhang, C. Y., Yin, L. J. & Dai, S. L. Diversity of root-associated fungal endophytes in *Rhododendron fortunei* in subtropical forests of China. *Mycorrhiza* **19**, 417–423 (2009).
38. Hambleton, S. & Currah, R. S. Fungal endophytes from the roots of alpine and boreal Ericaceae. *Can. J. Bot.* **75**, 1570–1581 (1997).
39. Tian, W., Zhang, C., Qiao, P. & Milne, R. Diversity of culturable ericoid mycorrhizal fungi of *Rhododendron decorum* in Yunnan, China. *Mycologia* **103**, 703–709 (2011).
40. Xiao, G. P. The role of root-associated fungi in the dominance of *Gaultheria shallon*. 1–148 PhD thesis, University of British Columbia, Vancouver (1994).
41. Berch, S. M., Allen, T. R. & Berbee, M. L. Molecular detection, community structure and phylogeny of ericoid mycorrhizal fungi. *Plant Soil* **244**, 55–66 (2002).
42. Wurzbürger, N., Higgins, B. P. & Hendrick, R. L. Ericoid mycorrhizal root fungi and their multicopper oxidases from a temperate forest shrub. *Ecol. Evol.* **2**, 65–79 (2011).
43. Allen, T. R., Millar, T., Berch, S. M. & Berbee, M. L. Culturing and direct DNA extraction find different fungi from the same ericoid mycorrhizal roots. *New Phytol.* **160**, 255–272 (2003).
44. Sun, L. F. *et al.* Different Distribution Patterns between Putative Ericoid Mycorrhizal and Other Fungal Assemblages in Roots of *Rhododendron decorum* in the Southwest of China. *Plos One* **7**, e49867 (2012).
45. Kamal, S. & Varma, A. *Microbiology of Extreme Soils*. 177–203 (Springer Berlin Heidelberg, 2008).
46. Gorman, N. R. & Starrett, M. C. Host range of a select isolate of the ericoid mycorrhizal fungus *Hymenoscyphus ericae*. *Hortscience* **38**, 1163–1166 (2003).
47. Bougoure, D. S. & Cairney, J. W. G. Assemblages of ericoid mycorrhizal and other root-associated fungi from *Epacris pulchella* (Ericaceae) as determined by culturing direct DNA extraction from roots. *Environ. Microbiol.* **7**, 819–827 (2005).
48. Selošce, M. A. *et al.* Sebaciales are common mycorrhizal associates of Ericaceae. *New Phytol.* **174**, 864–878 (2007).
49. Vohník, M. & Albrechtová, J. The co-occurrence and morphological continuum between ericoid mycorrhiza and dark septate endophytes in roots of six European *Rhododendron* species. *Folia Geobot.* **46**, 373–386 (2011).
50. Grünig, C. R., Queloz, V., Sieber, T. N. & Holdenrieder, O. Dark septate endophytes (DSE) of the *Phialocephala fortinii* s.l.-Acephala applanata species complex in tree roots: classification, population biology, and ecology. *Botany* **86**, 1355–1369 (2008).
51. Ahlich-Schlegel, K. Vorkommen und Charakterisierung von dunklen, septierten Hyphomyceten (DSH) in Gehölzwurzeln. PhD thesis, Swiss Federal Institute of Technology, Zürich (1997).
52. Brundrett, M. C. Understanding the roles of multifunctional mycorrhizal and endophytic fungi. *Microbial root endophytes* 281–298 (Springer Berlin Heidelberg, 2006).
53. Piercey, M., Thormann, M. & Currah, R. Saprobic characteristics of three fungal taxa from ericacean roots and their association with the roots of *Rhododendron groenlandicum* and *Picea mariana* in culture. *Mycorrhiza* **12**, 175–180 (2002).
54. Read, D. J. The structure and function of the ericoid mycorrhizal root. *Ann. Bot.* **77**, 365–374 (1996).
55. Bergero, R., Girlanda, M., Bello, F., Luppi, A. & Perotto, S. Soil persistence and biodiversity of ericoid mycorrhizal fungi in the absence of the host plant in a Mediterranean ecosystem. *Mycorrhiza* **13**, 69–75 (2003).
56. Toju, H., Sato, H., Yamamoto, S., Kadowaki, K. & Tanabe, A. S. How are plant and fungal communities linked to each other in belowground ecosystems? A massively parallel pyrosequencing analysis of the association specificity of root-associated fungi and their host plants. *Ecol. Evol.* **3**, 3112–3124 (2013).
57. Davey, M. L., Heegaard, E., Halvorsen, R., Ohlson, M. & Kausarud, H. Seasonal trends in the biomass and structure of bryophyte-associated fungal communities explored by 454 pyrosequencing. *New Phytol.* **195**, 844–856 (2012).
58. Beeck, M. O., Ruytinx, J., Smits, M. M., Vangronsveld, J. & Colpaert, J. V. Belowground fungal communities in pioneer Scots pine stands growing on heavy metal polluted and non-polluted soils. *Soil Biol. Biochem.* **86**, 58–66 (2015).
59. Oja, J., Kohout, P., Tedersoo, L., Kull, T. & Koljalg, U. Temporal patterns of orchid mycorrhizal fungi in meadows and forests as revealed by 454 pyrosequencing. *New Phytol.* **205**, 1608–1618 (2015).
60. Buscardo, E. *et al.* Contrasting soil fungal communities in Mediterranean pine forests subjected to different wildfire frequencies. *Fungal Divers.* **70**, 85–99 (2015).
61. Begerow, D., Nilsson, H., Unterseher, M. & Maier, W. Current state and perspectives of fungal DNA barcoding and rapid identification procedures. *Appl. Microbiol. Biotechnol.* **87**, 99–108 (2010).
62. Ekblom, R. & Galindo, J. Applications of next generation sequencing in molecular ecology of non-model organisms. *Heredity* **107**, 1–15 (2011).
63. Lindahl, B. D. *et al.* Fungal community analysis by high-throughput sequencing of amplified markers—a user’s guide. *New Phytol.* **199**, 288–299 (2013).

64. Buée, M., Vairelles, D. & Garbaye, J. Year-round monitoring of diversity and potential metabolic activity of the ectomycorrhizal community in a beech (*Fagus sylvatica*) forest subjected to two thinning regimes. *Mycorrhiza* **15**, 235–245 (2005).
65. Schweitzer, J. A. *et al.* Forest gene diversity is correlated with the composition and function of soil microbial communities. *Popul. Ecol.* **53**, 35–46 (2011)
66. Toberman, H., Freeman, C., Evans, C., Fenner, N. & Artz, R. R. Summer drought decreases soil fungal diversity and associated phenol oxidase activity in upland *Calluna* heathland soil. *FEMS Microbiol. Ecol.* **66**, 426–436 (2008).
67. Fujimura, K. E. & Egger, K. N. Host plant and environment influence community assembly of High Arctic root-associated fungal communities. *Fungal Ecol.* **5**, 409–418 (2012).
68. Day, M. J. & Currah, R. S. Role of selected dark septate endophyte species and other hyphomycetes as saprobes on moss gametophytes. *Botany* **89**, 349–359 (2011).
69. Grunewaldt-Stöcker, G., Von den Berg, C., Knopp, J. & Von Alten, H. Interactions of ericoid mycorrhizal fungi and root pathogens in Rhododendron: *In vitro* tests with plantlets in sterile liquid culture. *Plant Root* **7**, 33–48 (2013).
70. Fichtner, A. *et al.* Effects of anthropogenic disturbances on soil microbial communities in oak forests persist for more than 100 years. *Soil Biol. Biochem.* **70**, 79–87 (2014).
71. Walker, C. Arbuscular mycorrhiza in the Living Collections at the Royal Botanic Garden Edinburgh. *Sibbaldia: the Journal of Botanic Garden Horticulture* **11**, 143–157 (2013).
72. Johnson, N. C., Angelard, C., Sanders, I. R. & Kiers, E. T. Predicting community and ecosystem outcomes of mycorrhizal responses to global change. *Ecol. Lett.* **16**, 140–153 (2013).
73. Bougoure, D. S., Parkin, P. I. & Cairney, J. W. G. Diversity of fungi in hair roots of Ericaceae varies along a vegetation gradient. *Mol. Ecol.* **16**, 4624–4636 (2007).
74. Glinka, C. & Hawkes, C. V. Environmental controls on fungal community composition and abundance over 3 years in native and degraded shrublands. *Microb. Ecol.* **68**, 807–817 (2015).
75. Zhang, L. W., Mi, X. C., Shao, H. B. & Ma, K. P. Strong plant-soil associations in a heterogeneous subtropical broad-leaved forest. *Plant Soil* **347**, 211–220 (2011).
76. Yu, M. J., Hu, Z. H., Yu, J. P., Ding, B. Y. & Fang, T. Forest vegetation types in Gutianshan Natural Reserve in Zhejiang. *J. Zhejiang Univ. Sci.* **27**, 375–380 (2001).
77. Song, K. *et al.* Variation in phylogenetic structure of forest communities along a human disturbance gradient in Gutianshan forest. *China. Biodivers. Sci.* **19**, 190–196 (in Chinese with English abstract) (2011).

Acknowledgements

We thank Dr. Xiaojuan Liu for providing climatic information of the sampling sites, and Yefei Jin and Li Han for preparing the material and experiments. This study was supported by the National Natural Science Foundation of China (31170469, 31170495 and 31470565) and Technology division of Shaoxing (2013B70040). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Conceived and designed the experiments: L.S., K.P. and Y.Z. Performed the experiments: Y.Z., J.N. and F.T. Analyzed the data: Y.L. and L.S. Contributed reagents/materials/analysis tools: L.S., Y.Z., N.J. and F.T. Wrote the paper: Y.Z., Y.L., L.J., L.S. and Y.Q.L.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, Y. *et al.* Root-associated fungi of *Vaccinium carlesii* in subtropical forests of China: intra- and inter-annual variability and impacts of human disturbances. *Sci. Rep.* **6**, 22399; doi: 10.1038/srep22399 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>